


Please rate each of the following measures according to the strength of your recommendation (1=Low; 2=Moderate; 3=Strong) that the measure be included in the SIM Quality Measure Set.

Answered: 16 Skipped: 0



	1 = Low	2 = Moderate	3 = Strong	Total	Weighted Average		25.00%	31.25%	43.75%		
Diabetes: Hemoglobin A1c Poor Control (>9%) (NQF 0059)	0.00% 0	6.25% 1	93.75% 15	16	2.94	Heart Failure: Beta-blocker therapy for left ventricular systolic dysfunction (NQF 0083)	4	5	7	16	2.19
Hypertension: Controlling high blood pressure (NQF 0018)	0.00% 0	12.50% 2	87.50% 14	16	2.88	Use of spirometry testing in the assessment and diagnosis of COPD (NQF 0057)	4	8	4	16	2.00
Diabetes eye exam (NQF 0055)	0.00% 0	31.25% 5	68.75% 11	16	2.69	Disease modifying anti-rheumatoid arthritis (NQF 0054)	6	9	1	16	1.69
Diabetes: Medical attention for nephropathy (NQF 0062)	12.50% 2	12.50% 2	75.00% 12	16	2.83	Cardiac stress imaging: Testing in asymptomatic low risk patients (NQF 0672)	8	5	3	16	1.69
Appropriate treatment for children with upper respiratory infection (NQF 0069)	0.00% 0	37.50% 6	62.50% 10	16	2.83	Cardiac stress imaging: Preoperative evaluation in low risk surgery patients (NQF 0670)	9	4	3	16	1.83
Medication Management for people with asthma (NQF 1799)	12.50% 2	31.25% 5	56.25% 9	16	2.44	CAD: Medication adherence (no longer NQF endorsed)	11	3	2	16	1.44
Use of imaging studies for low back pain (NQF 0052)	0.00% 0	56.25% 9	43.75% 7	16	2.44						
Diabetes foot exam (NQF 0056)	12.50% 2	37.50% 6	50.00% 8	16	2.38						
CAD: Persistence of Beta blocker therapy after a heart attack (NQF 0071)	6.25% 1	50.00% 8	43.75% 7	16	2.38						
Avoidance of antibiotic treatment in adults with acute bronchitis (NQF 0058)	12.50% 2	37.50% 6	50.00% 8	16	2.38						
Ischemic vascular disease: use of aspirin or another anti thrombotic (NQF 0068)	25.00% 4	18.75% 3	56.25% 9	16	2.31						